

RELATION BETWEEN ANTIOXIDANT ACTIVITY OF AFFINED C SUGAR AND THERMALY INDUCED NONENZYMATIC BROWNING

Aleksandra Č. Mišan¹, Julianna F. Gyura², Marijana B. Sakač¹, Dragana M. Simović-Šoronja², Zita I. Šereš², Biljana S. Pajin²

¹Institute of Food Technology, University of Novi Sad, Bul. cara Lazara 1, 21000 Novi Sad, Serbia

²Faculty of Technology, University of Novi Sad, Bul. cara Lazara 1, 21000 Novi Sad, Serbia

* E-mail correspondence: aleksandra.misan@fins.uns.ac.rs

ABSTRACT: Affined C sugar was obtained at the third stage of crystallization from the process of sugar production and was characterized by 99.82% dry matter (DM) content, 97.6 %Z sucrose content, pH=6.81, 20366 IJ/DM coloured matter content measured at 420 nm and 4673 IJ/DM coloured matter content measured at 560 nm. As the development of coloured compounds and Maillard reaction products is favoured by the increase in temperature, the aim of this study was to investigate the relation between thermally induced nonenzymatic browning and antioxidant activity of affined C sugar. Nonenzymatic browning was tested by incubating 50 g of 40°Bx affined C sugar solution at 70 °C, 80 °C and 90 °C for 1200 min. Thermal treatment resulted in the increase in coloured substances content. Incubation at 90 °C for 1200 min resulted in partial sucrose degradation, which was followed by the increase in fructose and glucose content. Scavenging activity against DPPH radicals (DPPH') decreased with the increase in the incubation temperature. Since the antioxidant activity of Maillard reaction products was reported, our results indicate that Maillard reaction may not have been the main mechanism of nonenzymatic browning. Further research needs to be done in order to characterize the antioxidants in affined C sugar and the products of browning reaction under the described experimental conditions.

Key words: *affined C sugar; antioxidant activity; nonenzymatic browning*

INTRODUCTION

Colour of the final product is a key quality measurement in sugar industry. While the nature of colouring substances in sugarcane processing was well studied and documented (Bento and Sá, 1998), colour formation in sugar beet manufacture is not much examined and understood. Because of that, it is not possible to predict the content of coloured substances in the final product on the basis of raw material chemical composition.

Coloured substances are formed during sugar beet processing as a result of pH changes, and thermal and autocatalytic

effects (Šereš et al., 2004; Gyura et al., 2005). These impurities are of high molecular masses, polymeric and with tendency to occlude within the sugar crystal (Coca et al., 2008). Maillard reaction and alkaline degradation of invert sugars have been considered to be the main mechanisms of colour formation during purification stage (Coca et al., 2008). Melanoidins as Maillard reaction products are formed by the reaction of monosaccharides and carbonyl compounds with amino acids and are recognized as acidic and polymeric compounds with a highly complicated structure (Cämmerer and Kroh, 1995; Cämmerer et al., 2002).




Figure 1. Scheme of the three-stage sugar beet crystallization process

According to Heitz (1995), up to 80% of colour in sugar beet juice is a result of alkaline degradation of hexoses (HADP). The production of coloured HADP takes place at the

common pH of a beet sugar factory (8–11) while the formation of degradation products mainly occurs in the purification step where temperature increases up to 85 °C and pH

risers up to strong basic values (11–12). The nature and structure of coloured HADP have not been elucidated but they are probably formed due to the extensive aldolisation of intermediate di-carbonyl compounds in alkaline solutions (de Bruijn, 1986). Coloured substances formed through sugar beet processing negatively influence the quality of white sugar. However, Maillard reaction compounds have been reported to possess antioxidant activity (Yanagimoto et al., 2002; Yanagimoto et al., 2004). Payet et al. (2005) reported the antioxidant activity of cane brown sugars, which was attributed to the presence of polyphenol compounds and Maillard reaction products. As the concentration of nitrogen compounds in sugar beet juices is higher than in cane sugar juices, Coca et al. (2004) assume that the formation of melanoidins in sugar beet processing is more intensive.

Affined C sugar is an intermediate product of the three-stage crystallization process in sugar beet processing (Figure 1). From the third stage of crystallization, affined C sugar is directed to the first crystallization stage, and in this way directly influences the quality of the final product – white sugar. Since the development of coloured compounds and Maillard reaction products is favoured by the increase in temperature, the aim of this study was to investigate the relation between thermally induced nonenzymatic browning and antioxidant activity of affined C sugar.

MATERIALS AND METHODS

Materials

Affined C sugar was obtained at the third stage of crystallization from the process of sugar production. The basic characteristics of the affined C sugar were determined according to the methods specified by Reinefeld and Schneider (1978).

Thermally induced nonenzymatic browning

Affined C sugar was dissolved in distilled water and diluted to 40°Bx dry matter (con-

trol). Nonenzymatic browning was tested by incubating 50 g of affined C sugar solution (40°Bx dry matter) at 70 °C, 80 °C and 90 °C in water bath with the precision of ± 0.02 °C for 1200 min.

HPLC determination of fructose, glucose, raffinose and sucrose

A high-performance liquid chromatographic (HPLC) method was applied to determine fructose, glucose, raffinose and sucrose content of affined C sugar solutions. The HPLC (HP 1090, Hewlett-Packard, USA) system consisted of a Zorbax Carbohydrate 4.6 $\times$ 250 mm, 5 μ m column (Agilent Technologies), a solvent system of acetone-trile/water, 70/25 (v/v), a flow rate of 1 mL/min, a refractive index detector operating in the range of 4×10^{-5} RIU, and at the temperature of 35 °C. Prior to injection, all samples were diluted in deionised water and filtered through a 0.45 μ m Rotilabo-Spritzenfilter CME (Carl Roth).

Scavenging activity against DPPH' of affined C sugar

Affined C sugar was dissolved in deionized water, in an appropriate manner to obtain a series of concentrations. Scavenging activity against the stable DPPH' (1,1-diphenyl-2-picryl-hydrazyl radicals) was determined spectrophotometrically following the procedure of Espin and others (2000). The IC₅₀ (mg/mL) was defined as the mass concentration of an antioxidant solution which was required to quench 50% of the initial DPPH' under the given experimental conditions. It was obtained by interpolation from linear regression analysis.

RESULTS AND DISCUSSION

Basic quality parameters of affined C sugar are presented in Table 1. The obtained values are as expected for that type of an intermediate product. The contents of sucrose, fructose, glucose and raffinose in control and thermally treated solutions of affined C sugar diluted to 40°Bx dry matter were presented in Table 2.

Table 1.

Basic quality parameters of affined C sugar

Dry matter (%)	99.82
Content of sucrose (°Z)	97.6
pH value	6.81
Content of coloured matter at 420 nm (IJ/DM)	20366
Content of coloured matter at 560 nm (IJ/DM)	4673

Table 2.

The content of sucrose, fructose, glucose and raffinose in thermally treated solutions of 40° Bx affined C sugar

Sample	Sucrose (mg/mL)	Fructose (mg/mL)	Glucose (mg/mL)	Raffinose (mg/mL)
C 40 Bx	471.9 ± 1.68	0.97 ± 0.11	1.01 ± 0.20	2.10 ± 0.25
C 40 Bx 70 °C	557.1 ± 1.24	1.75 ± 0.18	2.23 ± 0.45	2.31 ± 0.18
C 40 Bx 80 °C	565.6 ± 1.32	1.86 ± 0.14	2.24 ± 0.32	2.59 ± 0.23
C 40 Bx 90 °C	549.1 ± 1.45	12.15 ± 0.16	15.52 ± 0.65	2.31 ± 0.24

Values are means of three determinations ± standard deviation.

Table 3.

Scavenging activity against DPPH' of thermally treated solutions of 40°Bx affined C sugar solution

Sample	DPPH' scavenging activity (IC₅₀, g/mL)
C sugar	0.440 ± 0.026
C sugar at 70 °C	0.644 ± 0.114
C sugar at 80 °C	1.613 ± 0.236
C sugar at 90 °C	4.010 ± 0.179

Values are means of three determinations ± standard deviation.

As it can be seen in Table 2, the content of sucrose, glucose and fructose slightly increased as a result of the incubation at 70 and 80 °C. Those results can be explained by the reduction of solvent volume during the incubation. However, 1200 min incubation at 90 °C resulted in partial sucrose degradation, which was followed by the increase in fructose and glucose content. Raffinose was present in all of the investigated solutions in the concentration which was not affected by the thermal treatment.

As the thermal treatment resulted in the increase in coloured substances content, which was measured as an increase in absorbance at 420 and 560 nm (data not shown), we decided to evaluate the antioxidant activity of thermally treated affined C sugar solutions. Scavenging activity against DPPH' test was chosen as it is one of the most frequently used and most convenient

tests. The results of scavenging activity against DPPH' of investigated solutions are presented in Table 3.

Development of Maillard reaction products is favoured by the increase in temperature, and considering that the antioxidant activity of Maillard reaction products was reported (Yanagimoto et al., 2002; Yanagimoto et al., 2004), the increase in antioxidant activity was expected. However, referring to the results shown in Table 3, scavenging activity on DPPH' decreased with the increase in the incubation temperature (higher IC₅₀ value indicates lower scavenging activity on DPPH'). The decrease in antioxidant activity, attributed to the action of Maillard reaction products, indicated that Maillard reaction may not have been the only mechanism of nonenzymatic browning. Further research needs to be done in order to characterize the antioxidants in affined C

sugar and the products of browning reaction under the described experimental conditions.

CONCLUSIONS

Prolonged incubation of affined C sugar solution at 90 °C induced sucrose degradation, which was followed by the increase in glucose and fructose content. Thermal treatment of affined C sugar favoured the formation of coloured substances and was, at the same time, followed by the decrease in antioxidant activity.

ACKNOWLEDGMENTS

This work was supported by the Project No TR-31014, Ministry of Education and Science, Republic of Serbia.

REFERENCES

1. Bento, L.S.M., Sá, S. (1998). Study of high molecular weight compounds in sugar using gel-permeation chromatography with an evaporative light scattering detector. *Carbohydrate Polymers*, 37, 257-261.
2. Cämmerer, B., Jalyschkov, V., Kroh, L. W. (2002). Carbohydrate structures as part of the melanoidin skeleton. *International Congress Series*, 1245, 269-273.
3. Cämmerer, B., Kroh, L.W. (1995). Investigations of the influence of reaction conditions on the elementary composition of melanoidins. *Food Chemistry*, 53, 55-59.
4. Coca, M., Garcia, M.T., González, G., Peña, M., Garcí, J.A. (2004). Study of coloured components formed in sugar beet processing. *Food Chemistry*, 86, 421-433.
5. Coca, M., García, M.T., Mato, S., Cartón, Á., González, G. (2008). Evolution of colorants in sugarbeet juices during decolorization using styrenic resins, *Journal of Food Engineering*, 89, 429-434.
6. De Bruijn, J. M. (1986). Monosaccharides in alkaline medium: isomerization, degradation and oligomerization. PhD Thesis, University of Technology, Delft, The Netherlands.
7. Espin, J. C., Soler-Rivas, C., Wichers, H. J. (2000). Characterization of the total free radical scavenger capacity of vegetable oils and oil fractions using 2,2-diphenyl-1-picrylhydrazyl radical. *Journal of Agricultural and Food Chemistry*, 48, 648-656.
8. Gyura, J., Šereš, Z., Eszterle, M. (2005). Influence of operating parameters on separation of green syrup coloured matter from sugar beet by ultra- and nano-filtration. *Journal of food engineering*, 66, 89-96.
9. Heitz, F. (1995). Les colorants en sucrerie. *Actualités Techniques et Industrielles*, 538-544.
10. Payet, B., Shum Cheong Sing, A., Smadja, J. (2005). Assessment of antioxidant activity of cane brown sugars by ABTS and DPPH radical scavenging assays: Determination of their polyphenolic and volatile constituents. *Journal of Agricultural and Food Chemistry*, 53, 10074-10079.
11. Reinefeld, E., Schneider, F. (Eds.) (1978). *Analitische Betriebskontrolle der Zuckerindustrie (Teil B)*, Verlag Dr. Albert Bartens, Berlin.
12. Šereš, Z., Gyura, J., Eszterle, M., Vatai, Gy. (2004). Coloured matters removal from sugar beet industry syrup by ultra- and nano-filtration. *Acta alimentaria*, 33, 119-127.
13. Yanagimoto, K., Lee, K.-G., Ochi, H., Shibamoto, T. (2002). Antioxidative activity of heterocyclic compounds found in coffee volatiles produced by Maillard reaction. *Journal of Agricultural and Food Chemistry*, 50, 5480-5484.
14. Yanagimoto, K., Ochi, H., Lee, K.-G., Shibamoto, T. (2004). Antioxidative activities of fractions obtained from brewed coffee. *Journal of Agricultural and Food Chemistry*, 52, 592-596.

ТЕРМИЧКИ ИНДУКОВАНО НЕЕНЗИМСКО ТАМЊЕЊЕ И АНТИОКСИДАТИВНА АКТИВНОСТ АФИНИСАНОГ Ц ШЕЋЕРА

Александра Ч. Мишан¹, Јулиана Ф. Ђура², Маријана Б. Сакач¹, Драгана М. Симовић-Шороња²,
Зита И. Шереш², Биљана С. Пајин²

¹Институт за прехранбене технологије, Универзитет у Новом Саду, Булевар цара Лазара 1, 21000 Нови Сад, Србија

²Технолошки факултет, Универзитет у Новом Саду, Булевар цара Лазара 1, 21000 Нови Сад, Србија

Сажетак: Афинисани Ц шећер, добијен у трећем степену кристализације технолошког поступка производње шећера из шећерне репе, карактерише 99,82% суве материје (SM), 97,6 °Z сахарозе, pH=6,81, 20366 IJ/SM бојених материја измерених на 420 nm и 4673 IJ/SM бојених материја измерених на 560 nm. С обзиром на то да је развој бојених материја и настајање производа Mallard-ове реакције фаворизовано са порастом температуре, циљ овога рада био је да се испита повезаност термички индукованог неензимског тамњења и антиоксидативне активности афинисаног Ц шећера. Неензимско тамњење је праћено инкубирањем 50 g раствора афинисаног шећера (40°Bx) на 70 °C, 80 °C и 90 °C током 1200 min. Термички третман је резултирао порастом садржаја бојених материја. Инкубација афинисаног Ц шећера на 90 °C током 1200 min довела је до делимичне деградације сахарозе, праћене порастом садржаја фруктозе и глукозе. Антирадикалска активност на DPPH радикале (DPPH') смањивала се са порастом температуре инкубације. Будући да је познато да производи Mallard-ове реакције испољавају антиоксидативну активност, добијени резултати указују да Mallard-ова реакција не представља главни механизам неензимског тамњења. Неопходна су даља истраживања како би се окарактерисале материје одговорне за антиоксидативну активност афинисаног Ц шећера, као и производи реакције неензимског тамњења генерисани у описаном експерименту.

Кључне речи: афинисани Ц шећер, антиоксидативна активност, неензимско тамњење

Received: 9 November 2011

Accepted: 30 November 2011